

Situation: Summing the Natural Numbers
Prepared at Penn State
Mid-Atlantic Center for Mathematics Teaching and Learning
14 July 2005 – Shari & Anna

Prompt

The course was a mathematical modeling course for prospective secondary mathematics teachers.

The discussion focused on finding an explicit formula for a sequence expressed recursively. During the discussion, a need to sum the integers from 1 to n was expressed. After several attempts to remember the formula, a student hypothesized that the formula contained n and $n + 1$. Another student said he thought it was $n(n + 1)/2$, but was not sure. During the ensuing discussion, a third student asked, "How do we know that $n(n + 1)/2$ is the sum of the integers from 1 to n , and won't this formula sometimes give fractions?"

Mathematical Foci

Mathematical Focus 1

Let S be the sum of the first n natural numbers. Then $S = 1 + 2 + 3 + \dots + n-1 + n$. By the commutative and associative properties of addition, $S + S = 1 + 2 + 3 + n-1 + n + 1 + 2 + 3 + \dots + n-1 + n = (n + 1) + (n-1 + 2) + (n-2 + 3) + \dots + (2 + n-1) + (1 + n) = n(n + 1)$. So $2S = n(n + 1)$, and $S = n(n + 1)/2$. Thus the sum of the first n natural numbers is $n(n + 1)/2$.


We know n and $n+1$ are consecutive integers, thus one is odd and the other even. Since the product of an even number and any other natural number is always even, our numerator is even, and thus divisible by two. Thus our formula always yields a natural number.

Mathematical Focus 2

We can represent the first n natural numbers with a triangular array of dots, with each row of dots representing a number. The following array represents the first four natural numbers.




If we make a copy of that array and rotate it so that, when joined with the original we obtain a rectangle, we can see that the dimensions of the rectangular array for this specific case are 4 by 5, and in the general case are n by $n + 1$.



Thus we have $4 \cdot 5 = 20$ dots in the rectangular array above ($n(n + 1)$ dots in the general case), and since we have two identical triangular arrays making up the rectangular array, there are $20/2 = 10$ dots in one triangular array ($n(n + 1)/2$ dots in one triangular array in the general case). Then since the triangular array represented the first n natural numbers, we have $n(n + 1)/2$ as the sum of the first n natural numbers. Since in every case we are working with two identical triangular arrays of dots representing natural numbers, it is clear that our formula will always yield a natural number.

A second way to visualize this is to use unit squares in place of the dots as seen below. The unit squares also form a triangular shape that can be copied and rotated 180 degrees to form a rectangle. One-half the area of this n by $n + 1$ rectangle is the sum of the first n natural numbers.



Mathematical Focus 3

We can represent the first n natural numbers with a triangular array of dots, with each row of dots representing a number. The following array represents the first four natural numbers.




One way to verify the formula would be to rearrange the dots into a rectangle. If n is even, we would have $n/2$ rows, as shown in the first figure below.



The remaining rows will be rotated and moved to complete a rectangle having dimensions $n/2$ by $n + 1$ as shown in the second figure above.

If n is odd, we would have $(n + 1)/2$ rows, as shown in the first figure below.



The remaining rows will be rotated and moved to complete a rectangle having dimensions $(n + 1)/2$ by n as shown in the second figure above.

Since this method only requires rearranging the dots, it is clear that it will always yield a natural number result.

Mathematical Focus 4

We can use mathematical induction to prove that our formula will always work. This proof does not derive the formula, but proves that the formula works once we have it.

Let sum be $\sum_{i=1}^n i$.

Our inductive hypothesis is that $S_n = \frac{n(n+1)}{2}$.

When $n = 1$, we have $1 = 1(2)/2 = 1$, which establishes our base case.

We assume that, for all numbers less than or equal to m , $S_m = \sum_{i=1}^m i = \frac{m(m+1)}{2}$. We

must show that this also holds for $m + 1$, that is, we must show that

$$\sum_{i=1}^{m+1} i = \frac{(m+1)(m+2)}{2}.$$

$$\sum_{i=1}^{m+1} i = \sum_{i=1}^m i + m + 1 = \frac{m(m+1)}{2} + m + 1 = \frac{m(m+1)}{2} + \frac{2(m+1)}{2} = \frac{(m+1)(m+2)}{2} \text{ as required.}$$

Thus the sum of the first n natural numbers is $n(n + 1)/2$.


We know n and $n+1$ are consecutive integers, thus one is odd and the other even. Since the product of an even number and any other is always even, our

numerator is even, and thus divisible by two. Thus our formula always yields a natural number.

Mathematical Focus 5

We know this is not a proof. Considering these questions could lead to an exploration of the sum of the first n natural numbers. For example, suppose that $n = 16$. One could begin by adding the numbers (i.e. $1 + 2 + 3 + \dots + 16$). However, one way to add the numbers, using the both the commutative and associative laws of addition, could be to change the order and groupings of the numbers. In our example, the first grouping could be the largest number with the smallest number (i.e. $1 + 16$), next grouping the second largest number with the second smallest number (i.e. $2 + 15$), then grouping the third largest number with the third smallest number ($3 + 14$), and so forth until all possible groupings are created. We will have 8 groups that sum to 17, since 8 is half of sixteen. Thus the sum of the numbers from 1 to 17 is $8 \cdot 17 = 136$. We see that this is in line with the given formula of $n(n+1)/2$. However, this only shows that the formula seems to work and give a natural number when adding an even number of numbers.

Now suppose that $n = 9$. One could begin by adding the numbers (i.e. $1 + 2 + 3 + \dots + 9$). However, we could also group the numbers as above using the following diagram.




Notice that since 9 is odd, 5 is not paired with another number. Thus we have 4 groups that sum to 10. Notice that 5 is half of our group sum of 10, so we have $4 \frac{1}{2}$ groups of 10. $4 \frac{1}{2}$ is half of 9, and 10 is $9 + 1$, so our formula $n(n + 1)/2$ seems to hold. When we are adding an odd number of numbers, grouping as above will always result in even sums since we add odd to odd and even to even. Even though we will always have to deal with half of a group sum (as in the 5 in our example above), that sum is even, so the result is a natural number. We have obtained a natural number result of the formula when adding an odd number of numbers.

Mathematical Focus 6

We can represent the first n natural numbers by unit squares arranged in n rows with each row of squares representing a number. The sum of the first n natural

numbers is equal to the number of squares in the region, which, since the squares are unit squares, is equal to the area of the region.



The area of this region can be found by finding the area of the right triangle with legs of length n and adding the area of each of the small triangles left out of this triangle (highlighted above). The area of the large right triangle is $n^2/2$, and the area of each of the n small triangles is $1/2$, so the combined area is $n^2/2 + n/2 = n(n + 1)/2$.

If n is odd, then n^2 is odd, so $n^2 + n$ is even and thus our formula yields a natural number. If n is even, then n^2 is even, so $n^2 + n$ is even and our formula once again yields a natural number.

Mathematical Focus 7 & Extension

Polya (1981) describes how Pascal solved this problem and extended it to the general case of finding the sum of the k^{th} powers of the first n natural numbers. The method depends on expanding $(n + 1)^{k+1}$.

Let S_1 be the sum of the first n natural numbers. To find S_1 , expand $(n + 1)^2$ and write out its values for several numbers to see the pattern:

$$(n + 1)^2 = n^2 + 2n + 1 \text{ so } (n + 1)^2 - n^2 = 2n + 1.$$

$$(n + 1)^2 - n^2 = 2n + 1$$

$$(1) \quad 2^2 - 1^2 = 2 \cdot 1 + 1$$

$$(2) \quad 3^2 - 2^2 = 2 \cdot 2 + 1$$

$$(3) \quad 4^2 - 3^2 = 2 \cdot 3 + 1$$

⋮

$$(n) \quad (n + 1)^2 - n^2 = 2n + 1$$

Then we can add all the equations (1) through (n) together. The sum of these equations will be $(n + 1)^2 - 1 = 2(S_1) + n$. Then

$$S_1 = \frac{(n + 1)^2 - n - 1}{2} = \frac{(n + 1)^2 - 1(n + 1)}{2} = \frac{(n + 1 - 1)(n + 1)}{2} = \frac{n(n + 1)}{2}.$$

To find the sum of the first n squares, S_2 , expand $(n + 1)^3$ and write out its values for several numbers to see the pattern:

$$(n + 1)^3 = n^3 + 3n^2 + 3n + 1, \text{ so } (n + 1)^3 - n^3 = 3n^2 + 3n + 1$$

$$(n + 1)^3 - n^3 = 3n^2 + 3n + 1$$

$$2^3 - 1^3 = 3 \cdot 1^2 + 3 \cdot 1 + 1$$

$$3^3 - 2^3 = 3 \cdot 2^2 + 3 \cdot 2 + 1$$

$$4^3 - 3^3 = 3 \cdot 3^2 + 3 \cdot 3 + 1$$

⋮

$$(n + 1)^3 - n^3 = 3n^2 + 3n + 1$$

Again add all the equations together. The sum of these equations will be

$(n + 1)^3 - 1 = 3 \cdot S_2 + 3S_1 + n$. Substituting for S_1 and solving for S_2 gives us

$$S_2 = \frac{(n + 1)^3 - 3 \frac{n(n + 1)}{2} - n - 1}{3} = \frac{2(n + 1)^3 - 3n(n + 1) - 2(n + 1)}{6} = \frac{(n + 1)[2(n + 1)^2 - 3n - 2]}{6}$$

$$= \frac{(n + 1)(2n^2 + n)}{6} = \frac{n(n + 1)(2n + 1)}{6}$$

This method can be extended to find the sum of the first n cubes, and then the first n fourth powers and so on, as long as the sums of all the previous powers have been found.

References

Polya, G. (1981). *Mathematical discovery: On understanding, learning, and teaching problem solving*. John Wiley & Sons: New York.

END OF VIGNETTE – END OF VIGNETTE – END OF VIGNETTE