

# MAC-CPTM Situations Project

## Situation 40: Powers

Prepared at Penn State  
Mid-Atlantic Center for Mathematics Teaching and Learning  
14 July 2005 – Tracy, Jana, Christa, Jim

Edited at University of Georgia  
Center for Proficiency in Teaching Mathematics  
19 May 2006 – Eileen Murray, Bob Allen  
20 February 2007 – Sarah Donaldson

Edited cross-site (MAC-CPTM)  
29 March 2007 – Bob Allen, Kathy Heid, Shiv Karunakaran

### **Prompt**

During an Algebra I lesson on exponents, the teacher asked the students to calculate positive integer powers of 2. A student asked the teacher, “We’ve found  $2^2$  and  $2^3$ . What about  $2^{2.5}$ ?”

### **Commentary**

The prompt centers on the extension of the domain of the exponent from integers to rational numbers. The foci explore the nature of exponents numerically, analytically, and graphically. The numerical representation comprised of integral values in Focus 1 suggests a pattern for a curve for an extended domain that is illustrated in the graphical representation in Focus 3. Although Foci 1 and 3 provide an estimate, the analytical treatment in Focus 2 generates an exact value. These foci help expand the concept of exponentiation beyond repeated multiplication to accommodate the use of rational exponents.

It is also useful to consider the fact that  $f(x) = a^x$  would behave differently for particular values of “a” and “x.” In the discussion below, “a” is assumed to be positive and “x” is a rational number. If “a” had been negative, the arguments would not apply. Furthermore, different mathematical discussions would be necessary if “x” had been irrational, transcendental, or complex.

### **Mathematical Foci**

### Mathematical Focus 1

One method for estimating the value of  $2^x$  where  $x \in \mathbb{R}^*$  uses linear interpolation and the properties of the function  $f$  with rule  $f(n)=2^n$ , where  $n \in \mathbb{N}^*$ .


One way to calculate an estimate of the value  $2^{2.5}$  is to use linear interpolation between the two known values,  $2^2$  and  $2^3$ . The use of linear interpolation for estimating function values assumes that the function is well-behaved in that interval. Using linear interpolation, then  $2^{2.5}$  is approximately 6. However, it is important to understand that the value for  $2^{2.5}$  will not be exactly halfway between  $2^2$  and  $2^3$ . Whether the value of  $2^{2.5}$  is greater than 6 or less than 6 is determined by the pattern of growth of the function.

The table below shows a pattern of increasing growth between successive values of  $2^x$  and illustrates that exponential growth is different from a linear pattern of growth. Specifically, nonlinearity implies that even though 2.5 is the average of 2 and 3,  $2^{2.5}$  will not be the arithmetic mean of 4 and 8.

$x$	$2^x$
0	1
1	2
2	4
3	8
4	16

Since the differences between the successive values of  $2^x$  are increasing, one can argue that linear approximations will overestimate the value of  $2^{2.5}$ . Therefore,  $2^{2.5}$  should be less than 6.

Visualizing this numerical information and preserving the suggested curvature is also helpful in seeing why it makes sense that the value of  $2^{2.5}$  will be closer to 4 than to 8.



Specifically, if we look at this graph, we can begin to think of a way to connect the points that preserves the rate of growth in the table of values. This pattern of points suggests a graph that is concave up. We can use this fact to conclude further that the value of  $2^{2.5}$  will be closer to 4 than to 8.

### Mathematical Focus 2

*By defining rational powers in terms of integral powers, the properties of integral exponents are extended to rational exponents.*

By definition,  $b^{m/n}$  is the number that when raised to the  $n$ th power given  $b^m$  as a result. So,  $b^{m/n} = \sqrt[n]{b^m}$  for  $m \in \mathbb{Q}^*$ ,  $n \in \mathbb{Q}^+$ , and  $b \in \mathbb{Q}^+$ . Specifically,  $2^{1/2} = \sqrt{2}$ . Using this representation with the properties of exponents, this quantity can be represented by  $2^2 \cdot 2^{0.5} = 2^2 \sqrt{2} \approx 4(1.414) = 5.656$ . So  $2^{2.5} \approx 5.656$ . Similarly,  $(2^5)^{1/2} = \sqrt{2^5} = \sqrt{32} \approx 5.656$  or  $(2^{1/2})^5 = (\sqrt{2})^5 \approx 1.414^5 \approx 5.656$ . To rewrite  $2^{2.5}$  in another form,  $(2^{1/2})^5 = (\sqrt{2})^5 = \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} = 4\sqrt{2}$ .

Consistency is maintained in this focus. The choices that are made in the extension of the definition and properties of exponents are made so that they are preserved in a different number system, i.e. from whole numbers to rational numbers. That is, we can define  $2^{\text{rational}}$  in a way that allows the properties of

exponents to hold. By deciding that the properties must hold, we can then use them to express  $2^{2.5}$  as  $(2^5)^{\frac{1}{2}}$  or  $(2^{\frac{1}{2}})^5$ .

### Mathematical Focus 3

A graphical representation of the function  $f$  with rule  $f(x) = 2^x$  allows for more accurate estimation of  $2^n$  where  $n \in \mathbb{R}$ .

It should be recognized that at this point one is using the assumption that the domain of the function  $f$  with rule  $f(x) = 2^x$  can be extended from the set of integers to the set of real numbers. In particular, the resulting graph of the function with the new domain will be represented by a smooth, continuous graph of the function. This graph allows us to obtain an estimate for  $f(2.5)$  with varying degrees of accuracy depending on the technology or method employed.

For example, we can estimate the value of  $f(2.5)$  from a calculator-generated graph of  $f$  and the trace option to obtain a value of  $f(2.5)$ . On the other hand, we can look at the intersection of the function graph with the vertical line  $x = 2.5$ .

