

MAC-CPTM Situations Project

Situation 21: Exponential Rules

Prepared at University of Georgia
Center for Proficiency in Teaching Mathematics
29 June 2005 -- Erik Tillema

Edited at University of Georgia
16 June 2006 -- Sarah Donaldson

Prompt

A teacher in an Algebra II college preparatory class puts the following problem on the board $x^m \cdot x^n = x^5$. She has just gotten done with her lesson which reviewed the rules for exponents. She tells her students to take out a piece of paper and write down as many answers for m and n that the students think will make the statement true. She promises the students extra credit if they write all of them down. After allowing the students a few minutes to write out answers to the problem, one student raises her hand and says aloud so the students around her can hear, "How can I ever get extra credit? I keep thinking of new one's that work." Several other students in the vicinity seem to agree.

What ideas or focal points might the teacher use to capitalize on the students comments?

Commentary:

This prompt provides a wonderful opportunity to help students deepen their understanding of exponent rules, as well as their understanding of continuous functions and the notion of infinite solutions. The following foci address each of these issues.

Mathematical Foci

Mathematical Focus 1: Understanding why $x^m \cdot x^n = x^{m+n}$

$$x^m \cdot x^n = x^{m+n}$$

WHY?

x^m means x multiplied by itself m times, or $(x_1x_2x_3\dots x_m)$

x^n means x multiplied by itself n times, or $(x_1x_2x_3\dots x_n)$

Therefore $x^m \cdot x^n = (x_1x_2x_3\dots x_m)(x_1x_2x_3\dots x_n) = (x_1x_2x_3\dots x_{(m+n)})$

(that is, x multiplied by itself m times, and n more times, for a total of $(m+n)$ times, or x^{m+n}).

Mathematical Focus 2: $m + n = 5$ as a function


In this Situation, the specific problem is $x^m \cdot x^n = x^5$. Using the exponent rule $x^m \cdot x^n = x^{m+n}$,

$$x^m \cdot x^n = x^5$$

$$x^{m+n} = x^5$$

$$m + n = 5$$

$m + n = 5$ can be considered as a function:



$$m = f(n) = 5 - n$$

The graph shows that this is a continuous function. That is, it forms an unbroken set of points. The fact that this graph is a line (rather than a discrete number of points) indicates that there are infinite solutions to the equation $m + n = 5$.

In a problem in which m and n are exponents, students may initially assume that the values of m and n are limited to positive integers. If this were the case, the only solutions would be:

$$\begin{aligned}
 x^0 \cdot x^5 &= x^5 \\
 x^1 \cdot x^4 &= x^5 \\
 x^2 \cdot x^3 &= x^5 \\
 x^3 \cdot x^2 &= x^5 \\
 x^4 \cdot x^1 &= x^5 \\
 x^5 \cdot x^0 &= x^5
 \end{aligned}$$

However, m and n are not limited by this restriction—they could be any two numbers whose sum is 5. A few examples are:

$$\begin{aligned}
 x^7 \cdot x^{-2} &= x^5 \\
 x^{1.5} \cdot x^{3.5} &= x^5
 \end{aligned}$$

but of course there are an infinite number of examples.

Mathematical Focus 3: Plotting m and n on a number line.

If $m + n = 5$, then the average (mean) of m and n is 2.5. That is,

$$m + n = 5$$

$$\frac{m + n}{2} = \frac{5}{2} = 2.5$$

The mean of two values is shown on a number line in this manner: A pair of values has a mean of 2.5 if each value is represented as a point that is equidistant from 2.5 on the number line. In other words, two values m and n are represented as endpoints of a segment whose midpoint is 2.5.

Each unique pair of endpoints represents a unique solution to $m + n = 5$. Since the number line extends infinitely in opposite directions, there are infinite pairs of points whose midpoint is 2.5, and therefore infinite solutions to $m + n = 5$.

